

*Full Length Research Paper*

# Efficient control of conidium germination, mycelial growth and early blight in tomato *in vitro* with essential oils under farm conditions

C. L. M. Abreu<sup>1,2</sup>, C. I. M. A. Feres<sup>2,4,6\*</sup>, E. L. Furtado<sup>2</sup>, L. C. Ming<sup>2</sup>, M. O. M. Marques<sup>3</sup>, R. S. Acációd, A. Riffel<sup>5</sup>, H. F. Goulart<sup>4</sup>, A. E. G. Santana<sup>4</sup>, F. S. Bernardes<sup>4</sup> and F. L. A. Câmara<sup>2</sup>

<sup>1</sup>Empresa Mato-grossense de Pesquisa, Assistência e Extensão Rural do Mato Grosso - EMPAER-MT, Rua 02, s/n, Edifício Ceres, 3º andar, CPA, 78.050-250, Cuiabá – Mato Grosso, Brazil.

<sup>2</sup>Universidade Estadual Paulista, Faculdade de Ciências Agrônômicas, Depto. de Produção Vegetal, Fazenda Experimental Lageado, Rua Josá Barbosa de Barros, 1780, C. Postal 237, 18610-307, Botucatu – São Paulo, Brazil.

<sup>3</sup>Instituto Agronômico, Centro de Pesquisa e Desenvolvimento de Recursos Genéticos Vegetais, Seção de Fitoquímica, C. Postal 28, 13001-970, Campinas - São Paulo, Brazil.

<sup>4</sup>Instituto de Química e Biotecnologia, Universidade Federal de Alagoas, 57092-970 Maceió, Alagoas, Brasil.

<sup>5</sup>Empresa Brasileira de Pesquisa Agropecuária (Embrapa)- Tabuleiros Costeiros - UEP Rio Largo/AL, BR - 104 Norte, km 85, s/n - Zona Rural - Rio Largo/AL CECA - UFAL (Centro de Ciências Agrárias da UFAL) - Campus Delza Gitai, Cep: 57.100-000, Brazil.

<sup>6</sup>Faculdades Integradas do Norte de Minas, Campus JK, Avenida Osmani Barbosa, nº 11111, Bairro JK, Montes Claros, Minas Gerais, Brazil.

Received 6 May, 2016; Accepted 15 September, 2016

The aim of this work was to find a natural product and efficient dose to control early blight in tomato crop in agro-ecological or conventional agriculture system. The treatments were essential oils from *Cymbopogon citratus*, *Eucalyptus citriodora*, *Piper hispidinervum*, *Cymbopogon martini*, *Rosmarinus officinalis*, *Syzygium aromaticum*, *Cinnamomum zeylanicum*, *Mentha piperita*, *Citrus sinensis* var. *dulcis*, and *Melaleuca alternifolia* used against *Alternaria solani* in *Solanum lycopersicum* under greenhouse conditions. *In vitro*, mycelial growth and conidia germination were completely inhibited by *C. zeylanicum*, *C. martini*, *C. citratus*, and *S. aromaticum* oils of 750  $\mu\text{L L}^{-1}$ , *E. citriodora* and *M. alternifolia* at 2000  $\mu\text{L L}^{-1}$ , and *Mentha piperita* in 5000  $\mu\text{L L}^{-1}$ . In tomato, plants under greenhouse conditions, early blight in leaf was inhibited by *C. zeylanicum*, *C. martini*, *C. citratus*, and *S. aromaticum* at 750  $\mu\text{L L}^{-1}$ , and *E. citriodora* oil of 5000  $\mu\text{L L}^{-1}$ . The best results for MIC 50 and 90 with minor dose to control was essential oil from *C. citratus* at 341.32 and 1.822.10  $\mu\text{L L}^{-1}$  rich in geranial (46.91%), neral (34.34%), and geranyl acetate (6.30%). Therefore, it is possible to use this essential oils or manufacture a new and efficient product to control *Alternaria solani* in tomato plants.

**Key words:** Agroecology, *Alternaria solani*, *Cymbopogon* sp., *Eucalyptus* sp., Neral, *Solanum lycopersicon*.

## INTRODUCTION

Plant diseases are caused by fungi, nematodes, bacteria, and viruses, among which fungi are the main pathogens, causing great yield losses in numerous important crops

(Huang et al., 2010). Natural sources from plants play a significant role in the prevention and control of these diseases. In addition, products of higher plants may lead

to the discovery of the source of antimicrobial agents with possible novel mechanisms of action (Hada, 2014). Tomato (*Solanum lycopersicon* Mill.) is one of the most economically important vegetables in the world (Kurozawa and Pavan, 2011, 2005; Filgueira, 2000). It is the host for a wide spectrum of pests and diseases and thus requires specific ecological, nutritional and phytosanitary conditions in cropping and/or field (Leite et al., 2003). At post harvesting conditions, it is a highly perishable vegetable with a short shelf-life and high susceptibility to fungal disease during prolonged storage (Ibrahim, 2014).

Early blight, which is caused by *Alternaria solani* (Ell. and Martin) Jones and Grouet, is one of the most important diseases affecting tomato plants, since it causes direct (fruits) and indirect (branches and leaves) damages (Yanar, 2011; Leite et al., 2003). Fungicides including mancozeb, chlorothalonil, thiophanate-methyl and copper are used to control this disease.

Losses of up to 75% of agrochemicals during application have been recorded for tomato crops. Nowadays, producers use 45 agrochemical applications during the cycle of this species. The indiscriminate use of agrochemicals in tomato plants, associated with the lack of knowledge by producers, results in the development and intensification of occupational diseases and environmental contamination besides exposure of consumers to such risk (Araújo et al., 2000; Kishore and Pande, 2007; Fawzi et al., 2009).

Natural products have great potential for the management of pests and diseases, among other utilizations from extracts and essential oils (Zanella, 2015). Historically, the origin of pyrethroids and carbamates is an example of the potential of natural products for phytosanitary uses.

In this context, essential oils from plants have great potential due to their intrinsic characteristics such as high bioactivity, great molecule diversity and volatility, and brief environmental persistence (Knaak and Fiuza, 2010; Lee et al., 2008). The mechanism of action of essential oils in live cells is related to cell membrane permeability alteration, enzyme synthesis inhibition and/or inactivation (Souza et al., 2005). Cruz et al. (2015) had significant results in essential oils from *Plectranthus amboinicus* leaves against *Fusarium solani* at 10 µL. In *Pyricularia grisea*, *Cymbopogon winterianus* has total mycelia growth inhibition with 10 µL (Perini et al., 2013).

According to Soylu et al. (2010) and Copping and Duke (2007), based on the activity and mechanism of action concerning natural products, they may be directly applied on plants for protection and have potential applicability in the establishment of programs for the development of new products as well as synthesis and/or semi-synthesis processes. Also, Pawar and Thaker (2007) studied the antifungal effect of essential oils from 75 plant species

against *Alternaria porri* and *Fusarium oxysporum* f.sp. *ciceris*. Yanar et al. (2011) studied the antifungal effect of essential oils from 27 plant extracts, and Sallam and Kamal (2012) studied six plants. These authors emphasized the potentiality and importance of researching these compounds as an alternative for synthetic phytosanitary products, considering their economic and environmental viability.

Several authors have reported the efficiency and wide spectrum of action of essential oils against insects, fungi and pathogenic and phytopathogenic bacteria, both *in vitro* and *in vivo* (Paes, 2012.). Thus, the present work aimed at advancing applied research by evaluating, through preventive applications, the effect of different doses of essential oils from 10 plant species on conidium germination and mycelial growth *in vitro* as well as their controlling action, *in vivo*, on the severity of early blight caused by *A. solani* in the leaves of *S. lycopersicum* Mill. cv. Sta. Clara cultivated under greenhouse conditions.

## MATERIALS AND METHODS

Since the present paper is composed of three different linked works, the result was adopted as eliminatory parameters of treatment efficiency for the subsequent studies.

The employed essential oils are of commercial origin and obtained through hydrodistillation. In all experiments, essential oils were homogenized with a 1:1 mixture of water and Tween 80 detergent. The used concentrations were calculated based on the essential oil.

### The essential oils analysis

Analysis of essential oil components was analyzed through Gas Chromatography, ubMass Molecular Biology and Phytochemistry, Agronomical Institute of Campinas, São Paulo State, Brazil.

A 2 mg oil sample was diluted in 1 ml ethyl acetate (HPLC grade), from which a 1 µL aliquot was injected. The analysis was carried out in a GC/MS - Shimadzu/QP-5000, equipped with DB-5 column (30 m x 0.25 mm x 0.25 µm) electron ionization (70 eV) and mass scan range from 30 to 300 Da. Helium was used as carrier gas at 1.0 ml/min flow, and injector at 240°C. The following program was used: 50°C (5') to 160°C, 3°C per minute, and 160 to 220°C, 10°C per minute. Using a split ratio of 35, the temperatures of the ion source and GC-MS interface were 200 and 230°C, respectively. Compounds were identified by comparing their mass spectra with the GC/MS spectral library. Kovats retention index was calculated and determined by comparing the data with those found in literature (Adams, 1995).

### Effect of essential oils at different doses on *A. solani* mycelial growth *in vitro* (Experiment 1)

Mycelia were obtained in the Mycology and Forest Phytopathology Laboratory, Department of Plant Production, College of Agronomical Sciences, São Paulo State University-UNESP,

\*Corresponding author. E-mail: c.iezid@gmail.com.

Botucatu, São Paulo State, Brazil.

The experiment was carried out in Petri plates. The plates with each treatment were organized in a completely randomized design of 10 × 6 factorial arrangement, with five replicates. The treatments were essential oils from the following species: *Cymbopogon citratus* (lemon grass), *Eucalyptus citriodora* (lemon eucalyptus), *Piper hispidinervum* (long pepper), *Cymbopogon martini* (palmarosa), *Rosmarinus officinalis* (rosemary), *Syzygium aromaticum* (clove), *Cinnamomum zeylanicum* (cinnamon), *Mentha piperita* (peppermint), *Citrus sinensis* var. *dulcis* (orange), and *Melaleuca alternifolia* (tea tree), at 5 different concentrations (0, 250, 500, 750, 1000 and 5000  $\mu\text{L L}^{-1}$ ), except for *M. alternifolia*, *M. piperita* and *Eucalyptus citriodora* oils, which were used at 0, 1000, 2000, 3000, 4000 and 5000  $\mu\text{L L}^{-1}$ . After autoclaving, treatments were added to PDA culture medium. Plates were inoculated with 0.5 cm diameter disks and kept in BOD chamber at 25°C in the dark.

The colony diameter (mm) was daily evaluated until the eighth day. The mean of the ratio between measures in two perpendicular directions considered the growth value. The growth rate over time was assessed through parallelism test among doses of each oil, in which angular coefficients and constants from the growth linear regression equations were calculated. The results were plotted and the fungi static effect was analyzed through intercept (angular coefficient from the progress equation by each oil and dose), in which, the lower value, the smaller will be the angular coefficient of the disease symptom progression rate, mycelial growth and conidia germination at each treatment. Thus, each angular coefficient from the linear regression analysis contrasted the Scott-Knott method ( $p \leq 0.0001$ ).

To obtain the maximum inhibitory concentrations (50 and 90%), differential and integral calculus was applied. Regression statistical test was performed using "F" test.

#### Effect of essential oils at different doses on *A. solani* conidium germination *in vitro* (Experiment 2)

Conidia were obtained at Laboratory of Mycology and Forest Phytopathology, College of Agronomic Sciences, UNESP, Botucatu, São Paulo State, Brazil.

The experiment was carried out in 2% Agar medium in water. The design is completely randomized in 5 × 5 factorial arrangement, with three replicates. Treatments consisted of five essential oils selected among those from Experiment 1 (*C. citratus*, *C. martini*, *S. aromaticum*, *C. zeylanicum*, and *E. citriodora*). Each oil was tested at five concentrations (0, 250, 500, 750, and 1000  $\mu\text{L L}^{-1}$ ). The culture medium containing the treatments was applied onto glass slides placed in Petri plates and inoculated with 10  $\mu\text{L}$  of a suspension containing  $10^3$   $\mu\text{L L}^{-1}$  *A. solani* conidia. The plates were kept in a Biological organism development chamber (BOD) for 24 h at 25°C in the absence of light.

Conidium germination was evaluated with the aid of an optical microscope. The conidium that presented germ tube extension higher than the analyzed conidium length was considered germinated (Neely, 1978), and the germination percentage was calculated. Due to the pattern of conidium germination in response to treatments, only analysis of variance and mean contrasts were done through the Scott-Knott method ( $p \leq 0.0001$ ), to the detriment of regression analysis.

#### Effect of essential oils at different doses on the control of early blight caused by *A. solani* in the leaves of *S. lycopersicon* Mill. cv. Sta. Clara under greenhouse cropping (Experiment 3)

*S. lycopersicon* Mill. cv. Santa Clara seedlings (30 days old) was transplanted to 13-L pots. The substrate consisted of a mixture of earth, sand and commercial organic substrate (1:1:1), in addition to

10 kg fertilizer formula 4-14-08 per  $\text{m}^3$  of the mixture. Plants were kept for 57 days after transplanting (DAT) in a high tunnel under greenhouse condition; they were covered with 150  $\mu$  "Agrofilme" and subjected to daily irrigation through micro-aspiration (5 mm flow). The experimental design was in randomized blocks in 5 × 4 factorial arrangement, with 6 replicates; 5 essential oils (*C. citratus*, *E. citriodora*, *C. martini*, *S. aromaticum*, and *C. zeylanicum*) were used at 4 concentrations (0, 500, 750, and 1000  $\mu\text{L L}^{-1}$ ), except for *E. citriodora* (0, 750, 1000 and 5000  $\mu\text{L L}^{-1}$ ).

The pathogen inoculation consisted of the application of a  $10^4$  *A. solani* conidia/mL suspension twice on consecutive days after the first application of treatments, which was repeated at 3-day intervals until 57 DAT. It was done with the aid of a handheld backpack sprayer and plastic blanket for isolation of plots. At 57 DAT, the symptom severity was evaluated in the leaf tissue. The evaluator received a visual accuracy training with the software DISTRAN (public domain) of diagrammatic scale of damaged leaf area symptoms (Grade 0 = 0; 1 = 1-25%; 2 = 26-50%; 3 = 51-75%; and 4 = 76-100% injured leaf area) (Azevedo, 1997). Nine leaves from basal, middle and apical part of the canopy in ten plants were evaluated; from each replicate we can calculate the index control of relative efficiency (EFCR). The EFCR was calculated in relation to each standard treatment per treatments and replications (Ming et al., 2012).

Data were subjected to regression analysis ( $p \leq 0.0001$ ) with square root equations and Minimal Inhibitory Concentration (MIC) of 50 and 90% of symptoms of plant disease was determined. We do this study only in greenhouse field experiment with better oils after *in vitro* conidium and mycelial experiments.

## RESULTS AND DISCUSSION

### Chemical composition of the essential oils and their potential use

The results of GC/MS analysis from the essential oils tested and the activity reported on literature for each one and/or the most important compound detected are presented. The majority of antimicrobial compounds found in essential oils are terpenoids and phenylpropenes with the most active being phenols, although some aldehydes and non-phenolic substances also present promising antimicrobial activity. The target site and mode of action of most essential oil components still under is not well understood, especially in yeast (Hyldgaard, 2012).

The GC/MS results from the essential oils used in this work indicated that: The *C. citratus* (lemon grass) essential oil has the major constituents of geranial (46.91%), neral (34.34%), geranyl acetate (6.30%), camphene (1.02%), 6-methyl-5-hepten-2-one (1.28%), linalool (0.82%), citronellal (0.25%), isomenthol (1.81%), nerol (0.45%), geraniol (3.52%), and  $\alpha$ -trans-bergamotene (0.85%).

*C. martini* (palmarosa) essential oil had major component of geraniol (86.98%), followed by geranyl acetate (9.03%), *trans*-ocimene (1.45%), *trans*-caryophyllene (0.63%), and geranial (0.41%). Geranial and neral aldehyde mixture constitutes citral, a potent bioactive substance against phytopathogens (Glamočlija et al., 2011).

*E. citriodora* (eucalyptus) essential oil main compound was citronellal (74.74%), followed by citronellol (6.26%), isopulegol (5.18%), iso-isopulegol (4.33%), geraniol (2.96%),  $\alpha$ -pinene (0.45%),  $\beta$ -pinene (0.68%), limonene (0.33%), 1,8-cineole (1.51%), citronellyl acetate (1.24%), geranyl acetate (0.28%), and *trans*-caryophyllene (0.72%).

Both *S. aromaticum* (clove) and *C. zeylanicum* (cinnamon) had the major compound, eugenol: it is a major constituent in clove essential oil, and its antimicrobial activity is linked to its ability to permeabilize the cell membrane and interact with proteins; a phenylpropanoid of high antibiotic efficiency, specially against fungi; it can be used in crop phytopathogen control, as well as in food conservation. The antifungal mode of action of eugenol needs further investigation, but it is known to depend on cell proliferation. Eugenol treatment altered cell membrane and cell wall structures of proliferating *S. cerevisiae* cells resulting in the release of cellular content (Hyldgaard et al., 2012; Faria et al., 2006; Souza et al., 2005).

*S. aromaticum* had higher eugenol content (82.55%), relative to that of *C. zeylanicum* (73.45%). It had less diversity of substances - *trans*-caryophyllene (11.98%), eugenyl acetate (3.13%),  $\alpha$ -humulene (1.74), and  $\alpha$ -copaene (0.59%) compared with those present in *C. zeylanicum* essential oil: *Cis*-caryophyllene (4.67%), linalool (3.58%), benzyl benzoate (2.92%), *cis*-cinnamyl acetate (2.11%), safrole (1.25%),  $\alpha$ -pinene (1.24%),  $\alpha$ -phellandrene (1.32%), *ortho*-cymene (1.11%),  $\beta$ -phellandrene (1.10%), camphene (0.40%),  $\beta$ -pinene (0.47%),  $\alpha$ -terpineol (0.36%), *trans*-cinnamaldehyde (0.77%), and  $\alpha$ -copaene (0.75%). However, Pawar and Thaker (2007) attributed the antifungal activity of *C. zeylanicum* essential oil to cinnamaldehydes.

*M. alternifolia* (tea tree) had 4-terpineol (46.74%) and gamma-terpinene (18.16%) as the main components; it also presented  $\alpha$ -terpinene (7.5%), 1,8-cineole (5.25%), *ortho*-cymene (4.38%),  $\alpha$ -terpineol (3.43%), terpinolene (2.84%),  $\alpha$ -pinene (2.32%), limonene (1.69%),  $\alpha$ -selinene (1.27%), aromadendrene (1.00%), gamma-cadinene (0.99%), myrcene (0.82%),  $\beta$ -pinene (0.80%), sabinene (0.41%), *trans*-caryophyllene (0.37%), and delta-3-carene (0.34%).

*P. hispidinervum* (long pepper) presented Safrole with major constituent (93.85%) in the essential oil; it had  $\alpha$ -pinene (0.35%), myrcene (0.18%), delta-3-carene (0.41%), *ortho*-cymene (0.14%), limonene (0.21%), *cis*-ocimene (0.35%), *trans*-ocimene (0.88%), terpinolene (2.37%), *trans*-caryophyllene (0.30%), bicyclogermacrene (0.62%), and pentadecane (0.25%). In this oil, the major compound, Safrole detected has high relevance to industrial and agriculture chemical use. It has approximately 90 to 94% of the essential oil compound produced by the plant; it is applied as synergistic element in the composition of insecticides and herbicides (Maia et al., 1987).

For *R. officinalis* (rosemary), the essential oil is mainly composed of 1,8-cineole (25.51%),  $\alpha$ -pinene (22.23%), camphor (18.64%), camphene (10.63%), limonene (5.51%),  $\beta$ -pinene (4.76%), borneol (2.48%), myrcene (2.39%), 3-octanone (0.44%), *ortho*-cymene (1.87%), bornyl acetate (1.80%),  $\alpha$ -terpineol (1.40%), *trans*-caryophyllene (1.15%), and linalool (0.92%). Rosemary oil, until, has no agriculture use. *M. piperita* (peppermint) essential oil main constituent was menthol (45.18%), followed by menthone (22.54%), 1,8-cineole (7.03%), meta-cresol acetate (6.39%), menthyl acetate (5.97%), neo-menthol (2.88%), limonene (1.79%), pulegone (1.55%), *trans*-caryophyllene (1.21%),  $\beta$ -pinene (1.12%), non-identified compounds (0.72%), gamma-terpinene (0.70%),  $\alpha$ -pinene (0.60%), gamma-muurolene (0.59%), isomenthol (0.47%), sabinene (0.38%), 3-octanol (0.25%), *ortho*-cymene (0.25%), *trans*- $\beta$ -ocimene (0.25%), and myrcene (0.12%).

For *Citrus sinensis* var. *dulcis* (orange) essential oil used in this work is limonene (96.86%); it also presents small quantities of myrcene (2.27%) and  $\alpha$ -pinene (0.52%).

Considering the effect attributed to the essential oil major compound, this relation is almost direct when the oil has little diversity of molecules. It is important to point out the major compounds of essential oils, since they may be responsible, at least in part, for the oil antifungal activity (Salamci et al., 2007). However, is not true for all the essential oils and/ or microorganisms.

For essential oils with great diversity of molecules that is well distributed, the synergistic effect hypothesis is probably the most suitable. In some cases, large proportions of oxygenated monoterpenes may guarantee the essential oil antifungal activity (Santana, 2015).

The mechanism of action of essential oils in live cells is related to cell membrane permeability alteration, and enzyme synthesis inhibition and/or inactivation (Souza et al., According to the parallelism test for the growth curves of the analyzed doses of essential oils, *R. officinalis*, *C. sinensis* v. *dulcis*, and *P. hispidinervum* did not significantly reduce mycelial growth rate, presenting thus the highest angular coefficients (Table 1).

However, it can be seen that there are oils at different concentrations where the growth remains stagnant when the intercept remains at the abscissas (Table 1 and Figure 1). In this case, there is only fungistatic action and not fungicide effect. Aside oil type or dose, regular application is needed. However, such characteristics concerning monoterpenes constitute a dubious quality for the applicability of essential oils in crop protection, since they are highly volatile and instable. Commercial applications of essential oils would benefit from deeper insight into the mode of action behind individual compounds, as this could facilitate the exploitation of, e.g., synergistic combinations with more powerful antimicrobial properties (Hyldgaard, 2012).

The antifungal activity of a given essential oil may be

**Table 1.** Effect of different essential oils and doses on *A. solani* mycelial growth rate *in vitro*, according to the parallelism test of angular coefficients from linear regression equations for mycelial growth in colony diameter (mm) over eight days.

Oils	Doses ( $\mu\text{L L}^{-1}$ )					Mean R <sup>2</sup>	
	0	250	500	750	1000		
Angular coefficients ( $\alpha$ )							
<i>R. officinalis</i>	10.47 <sup>Aa</sup>	9.29 <sup>Aa</sup>	9.63 <sup>Aa</sup>	9.59 <sup>Aa</sup>	10.02 <sup>Aa</sup>	0.94	
<i>S. aromaticum</i>	10.47 <sup>Aa</sup>	7.46 <sup>Cb</sup>	6.37 <sup>Cc</sup>	0.00 <sup>Dd</sup>	0.00 <sup>Dd</sup>	0.96	
<i>C. citratus</i>	10.47 <sup>Aa</sup>	8.44 <sup>Bb</sup>	6.30 <sup>Cc</sup>	0.00 <sup>Dd</sup>	0.00 <sup>Dd</sup>	0.94	
<i>C. martini</i>	10.47 <sup>Aa</sup>	8.86 <sup>Bb</sup>	6.00 <sup>Cc</sup>	0.00 <sup>Dd</sup>	0.00 <sup>Dd</sup>	0.96	
<i>C. zeylanicum</i>	10.47 <sup>Aa</sup>	7.36 <sup>Cb</sup>	2.91 <sup>Dc</sup>	0.00 <sup>Dd</sup>	0.00 <sup>Dd</sup>	0.94	
<i>C. sinensis v. dulcis</i>	10.47 <sup>Aa</sup>	9.91 <sup>Aa</sup>	8.77 <sup>Bb</sup>	9.97 <sup>Aa</sup>	9.73 <sup>Aa</sup>	0.96	
<i>P. hispidinervum</i>	10.47 <sup>Aa</sup>	8.63 <sup>Bc</sup>	8.42 <sup>Bc</sup>	9.39 <sup>Ab</sup>	8.18 <sup>Bc</sup>	0.97	
	0	1000	2000	3000	4000	5000	
Angular coefficients ( $\alpha$ )							
<i>E. citriodora</i>	10.47 <sup>Aa</sup>	9.34 <sup>Ab</sup>	0.00 <sup>Bc</sup>	0.00 <sup>Bc</sup>	0.00 <sup>Bc</sup>	0.00 <sup>Ac</sup>	0.95
<i>M. piperita</i>	10.47 <sup>Aa</sup>	6.01 <sup>Bb</sup>	4.09 <sup>Ac</sup>	2.96 <sup>Ad</sup>	2.47 <sup>Ad</sup>	0.00 <sup>Ae</sup>	0.95
<i>M. alternifolia</i>	10.47 <sup>Aa</sup>	6.43 <sup>Bb</sup>	0.00 <sup>Bc</sup>	0.00 <sup>Bc</sup>	0.00 <sup>Bc</sup>	0.00 <sup>Ac</sup>	0.94

Means followed by the same uppercase letters in columns and lowercase letters in lines do not differ by the Scott-Knott test ( $p \leq 0.0001$ ).

isolated effects or synergic action by two or more compounds (Cruz et al., 2015). However, increasing amounts of evidence indicate that the inherent activity of essential oils may not rely exclusively on the ratio in which the main active constituents are present, but also interactions between these and minor constituents in the oils. Various synergistic antimicrobial activities have been reported for constituents or fractions of essential oils when tested in binary or ternary combinations (García-García et al., 2011). However, if we can use this technology at this moment, there is need to think about environment and economic sustainability questions under agroecology system prerogatives. In this case, the direct use of the essential oil on the cropping process to control insects and pathogenic microorganisms is possible. To the resources, free input given by farmers is possible in planting economic or alimentary plants with phytosanitary plants for essential oil production or other environmental services.

### Effect of essential oils at different doses on *A. solani* mycelial growth *in vitro* (Experiment 1)

*A. solani* mycelial growth was differently influenced, when the studied essential oils and their concentrations was compared (Table 1 and Figure 1).

Action of the mycelial growth control can be possible because it has a high variation on the essential oils compounds. There are two hypotheses for the complex action: First, synergic action among the compounds against *A. solani*; second, horizontal action with multiples attack points in the membranes and biochemical process

in the mycelial tissue.

In addition, Daferera et al. (2003) reported that *R. officinalis* oil was inefficient against *B. cinerea* and *F. solani* mycelial growth, corroborating the present results.







As such, essential oils did not show any action against *A. solani* under the conditions of the present work. They were eliminated at this stage. The opposite was observed for *E. citriodora*, *M. alternifolia*, *M. piperita*, *S. aromaticum*, *C. citratus*, *C. martini*, and *C. zeylanicum* oils, which not only reduced the colony growth rate but also completely inhibited it at the highest doses (Table 1).

For the remaining oils, the fungistatic effect for five days on colony stagnation was confirmed, and mycelial growth was completely inhibited at doses higher than 750 and 1000  $\mu\text{L L}^{-1}$  (Figure 1).


Similar to the results obtained in the present work, several authors have reported the fungicide and fungistatic effects of different doses of *S. aromaticum*, *C. zeylanicum*, *C. citratus*, *C. martini*, *E. citriodora*, *M. alternifolia* and *M. piperita* essential oils against *in vitro* pathogens development: *Aspergillus parasiticus*, *Aspergillus niger*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Rhizopus* sp, *Penicillium* sp, *Euopium repens*, *Rhizoctonia solani*, *Rhizoctonia bataticola*, *Fusarium moniliforme*, *F. oxysporum*, *F. solani*, *Bipolaris sorokiniana*, *Botrytis cinerea*, *Helminthosporium oryzae*, *Alternaria* sp., *Alternaria porri*, *Myrothecium verrucaria*, and *Curvalaria lunata* (Pawar and Thaker, 2007; Negrelle and Gomes, 2007; Faria et al., 2006; Fonseca et al., 2006).

Besides, Faria et al. (2006) reported *Alternaria* sp. was susceptible to the essential oil of *Ocimum gratissimum* chemotype-eugenol as well as to isolated eugenol. Pawar

Intercept







**Figure 1.** Fungistatic effect of essential oils and their concentrations on *A. solani* mycelium growth *in vitro*, on the axis “x” days for return to growth and “Y” axis angular growth rates. Intercept analysis between angular coefficients and constants of linear equations on mycelia growth.

and Thaker (2007) also observed that essential oils of other plant species rich in eugenol were efficient against *Alternaria porri*. Salgado et al. (2003) studied the bioactivity of several *Eucalyptus* sp. essential oils; *E. urophylla* oil was the most efficient; *E. citriodora* oil was against *F. oxysporum* Schlecht., *Botrytis sorokiniana* Shoemaker, and *Botrytis cinerea* Pers, with dose-dependent inhibition point.

The antifungal effect observed for *S. aromaticum* and *C. zeylanicum* is probably due to eugenol, the main compound in both species: 82.55 and 73.45%, respectively, of their composition. In addition, there are reports of the antifungal effect of isolated eugenol against *A. solani* (Faria et al., 2006), *Fusarium* spp., *Rhizopus* spp., and others (Souza et al., 2005).

Campaniello et al. (2010) found that eugenol (from 100

to 150  $\mu\text{g mL}^{-1}$ ) is an effective antifungal compound against *Aspergillus*, *Penicillium*, *Emmericella* and *Fusarium* spp., suggesting that this activity could be attributed, in part, to the presence of a phenolic group.

*C. citratus* oil antifungal effect may be related to its major compounds, neral (34.34%) and geranial (46.91%), or to their associate action with the remaining oil components. This hypothesis is confirmed by Negrelle and Gomes (2007), who reported that the mixture of these aldehydes constitutes citral, to which important antifungal activity is attributed. Souza et al. (2005) also detected potent antifungal activity by citral against some phytopathogens like *Fusarium* spp. and *Rhizopus* spp. The activity of *E. citriodora* oil was probably due to its main compound, citronellal (74%). This hypothesis supports the results obtained by Salamci et al. (2007),

**Table 2.** Effect of different essential oils and their doses on *A. solani* conidium germination percentage *in vitro*.

Oils	Doses ( $\mu\text{L L}^{-1}$ )					
	0	250	500	750	1000	5000
	Germinated conidia (%)					
<i>S. aromaticum</i>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>
<i>C. citratus</i>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>
<i>C. martini</i>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>
<i>C. zeylanicum</i>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>	0.0 <sup>Ab</sup>
<i>E. citriodora</i>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	100.0 <sup>Aa</sup>	0.0 <sup>Ab</sup>

Means followed by the same uppercase letters in columns and lowercase letters in lines do not differ by the Scott-Knott test ( $p \leq 0.0001$ ).

who stated that the antifungal activity of essential oils is generally attributed to their major compound. It must be considered, however, that in certain cases the antifungal action, as well as its intensity, may be a result of the synergistic effect among the oil constituents. For example, neral and/or geranial activities are potentiated when they are associated with myrcene (Onawunmi et al., 1984).

#### Effect of essential oils at different doses on *A. solani* conidium germination *in vitro* (Experiment 2)

The effects observed for conidium germination was similar to those observed for *A. solani* mycelial growth (Table 2).

Corroborating the present results, Caccioni and Guizzardi (1994) also detected an inhibitory effect by the same concentrations of *C. zeylanicum* and *Cymbopogon* sp essential oils on *Monilinia laxa*, *Mucor piriformis* and *Rhizopus stolonifer* mycelial growth and spore germination. *S. aromaticum*, *C. citratus*, *C. martini*, and *C. zeylanicum* essential oils similarly inhibited conidium germination at 750  $\mu\text{L L}^{-1}$ , whereas inhibition by *E. citriodora* oil occurred only at 5000  $\mu\text{L L}^{-1}$  (Table 2).

Mishra and Dubey (1994) observed conidia germination and mycelial growth inhibition in *F. moniliforme*, *A. flavus* and *A. fumigatus* and other 47 species under *C. citratus* essential oil. On the other hand, these authors explain the importance of the cropping seasons of *C. citratus* on the essential oil compounds and their effect on fungi control.

#### Effect of essential oils at different doses on the control of early blight caused by *A. solani* in the leaves of *S. lycopersicum* (*L. esculentum*) Mill. cv. Sta. Clara cultivated under greenhouse conditions (Experiment 3)

Supporting the results obtained in the *in vitro* experiments

for mycelial growth (Table 1 and Figure 1) and conidium germination (Table 2), *in vivo* studies indicated a significant reduction in early blight severity in the leaves of *S. lycopersicum* cv. Sta. Clara cultivated under field greenhouse conditions (Figures 2 and 3). However, we can identify the significant dose effect difference by the MIC 50, 90% and Ymax estimated by biological and mathematical model to reduce symptoms by each oil (Table 3).

*C. zeylanicum* essential oil reached its maximum response (78, 2%) lower than the remaining oils, indicating that even a ten-fold increase in its dose is inefficient (Figure 2, Table 3).

Only *E. citriodora* essential oil at 4.802,64  $\mu\text{L L}^{-1}$  and *C. citratus* oil at 3.172,80  $\mu\text{L L}^{-1}$  was 100% efficient in controlling early blight severity, relative to the remaining species. *S. aromaticum* and *C. martini* oils at 2.998,39  $\mu\text{L L}^{-1}$  had 92% and 2898.642  $\mu\text{L L}^{-1}$  90% control, respectively (Figures 2 and 3; Table 3).


The chemical compounds observed at this treatment was geranial (46.91%), neral (34.34%), geranyl acetate (6.30%), camphene (1.02%), 6-methyl-5-hepten-2-one (1.28%), linalool (0.82%), citronellal (0.25%), isomenthol (1.81%), nerol (0.45%), geraniol (3.52%), and  $\alpha$ -trans-bergamotene (0.85%) (Figure 4).

Nashwa et al. (2012) found that different concentrations of six plant extracts, *O. basilicum*, *A. indica*, *E. chamadulonsis*, *D. stramonium*, *N. oleander*, and *A. sativum*, significantly reduced the early blight disease. The best results for MIC 50 and 90 with minor dose necessary to control was essential oil from *C. citratus* at 341, 32 and 1.822,10  $\mu\text{L L}^{-1}$  respectively (Table 3).


These results are corroborated by Glamočlija et al. (2011) and Geromini et al. (2015) on the effectiveness of Geranial and Neral aldehyde mixture that constitutes citral, a potent bioactive substance that fights against phytopathogens, especially fungi and post-harvest and storage diseases.

Farming under agroecological system is possible with the use of these essential oils. If the farmer has this species in their land, they can proceed to the oil





**Figure 2.** Efficiency of essential oils in controlling *A. solani* symptom severity in *S. lycopersicum* (*L. esculentum*) leaf area at 57 days after transplant under greenhouse conditions. Tested concentrations: 0, 500, 750 and 5000  $\mu\text{L L}^{-1}$ . Equations tested by "F" method ( $p \leq 0.0001$ ).




**Figure 3.** Efficiency of *E. citriodora* essential oil in controlling *A. solani* symptom severity in *S. lycopersicum* (*L. esculentum*) leaf area at 57 days after transplant under greenhouse conditions. Tested concentrations: 0, 750, 1000, and 5000  $\mu\text{L L}^{-1}$ . Equations tested by "F" method ( $p \leq 0.0001$ ).

**Table 3.** Inhibitory concentration (MIC 50, 90 and maximum effect) of different essential oils and their on *A. solani* symptoms reduction in tomato plants at experimental under field greenhouse conditions from response models at Figures 2 and 3.

Oils	Inhibitory concentration per Reduction in symptomatic leaf area (%)			
	MIC 50 ( $\mu\text{L L}^{-1}$ )	MIC 90 ( $\mu\text{L L}^{-1}$ )	MIC. Max. ( $\mu\text{L L}^{-1}$ )	$\hat{Y}$ Max. (%)
<i>S. aromaticum</i>	309.86	2.172.86	2.998.39	92
<i>C. citratus</i>	341.32	1.822.10	3.172.80	100
<i>C. martini</i>	312.78	2.440.25	2.898.64	90
<i>C. zeylanicum</i>	413.50	*ni	2.613.71	78
<i>E. citriodora</i>	410.05	2.253.39	4.802.64	100

\*ni = oil is not efficient at this level; Equations tested by "F" method ( $p \leq 0.0001$ ).



**Figure 4.** Chemical structure by compounds detected on *C. citratus* oil used on this experiment. GC/MS and Chemdraw Prime 15.0.

extraction by using steam distillation, and MIC 90 with a spreader-sticker. However, two applications are needed per week. Moreover, it is now also possible to develop a new product based on natural compound with more studies around formulation and chemical stability on field.

## Conclusion

According to the results obtained in the present experiments, technological studies are recommended for the development of new phytosanitary products based on essential oils of *C. citratus*, and *E. citriodora* and possibly *S. aromaticum* and *C. martini* for the control of *A. solani* in tomato crop.

## Conflict of Interests

The authors have not declared any conflict of interests.

## ACKNOWLEDGEMENTS

The authors thank FAPESP for financial support, project n. 03/1332-5; CNPq for Doctorate scholarship to the first and second authors; DPV-Horticulture/FCA-UNESP; CNPq/PDJ Process n. 151200/2014-9.

## REFERENCES

- Adams RP (1995). Identification of Essential Oil Components by Gas Chromatography. Mass Spectrom. pp. 237-243.
- Araújo AC, Nogueira DP, Augusto LG (2000). Impacto dos praguicidas na saúde: estudo da cultura de tomate. Rev. Saúde Pública, 34(3):309-313.
- Azevedo LAS (1997). Manual de quantificação de doenças em plantas. São Paulo. 144 p.
- Caccione DRL, Guizzardi M (1994). Inhibition of germination and growth of fruit and vegetable postharvest pathogenic fungi by essential oil components. J. Essential Oil Res. Camberra, 6(2):173-179.
- Campaniello D, Sinigaglia M (2010). Antifungal activity of eugenol against *Penicillium*, *Aspergillus*, and *Fusarium* species. J. Food Prot. 73:1124-1128.
- Copping LG, Duke SO (2007). Natural products that have been used commercially as crop protection agents. Pest Manage. Sci. 63(6):524-554.
- Cruz TP, Alves FR, Mendonça RF, Costa AV, Jesus JWC, Pinheiro PF, Marins AK (2015). Fungicide activity of essential oil *Cymbopogon winterianus* jowit (citronella) against *Fusarium solani*. Biosci. J. Uberlandia 31(1):1-8.
- Daferera DJ, Ziogas BN, Polissiou MG (2003). The effectiveness of plant essential oils on the growth of *Botrytis cinerea*, *Fusarium* sp and *Clavibacter michiganensis* subsp. *michiganensis*. Crop Prot. 22:39-44.
- Faria TJ, Ferreira RS, Yassumoto L, Pinto De Souza JR, Ishikawa NK, Barbosa AM (2006). Antifungal Activity of Essential Oil Isolated from *Ocimum gratissimum* L. (eugenol chemotype) against Phytopathogenic Fungi. Braz. Arch. Biol. Technol. 49(6):867-871.
- Fawzi EM, Khalil AA, Afifi AF (2009). Antifungal effect of some plant extracts on *Alternaria alternata* and *Fusarium oxysporum*. Afr. J. Biotechnol. 8(11):2590-2597.
- Figueira FAR (2000). Novo manual de olericultura; Agrotecnologia moderna na produção e comercialização de hortaliças. 20 ed. Viçosa: UFV. P 402.
- Fonseca EN, Figer A, Furtado DT, Lopes D, Alviano DS, Alviano CS, Leitão SG (2006). Análise química e atividade antimicrobiana do óleo essencial dos frutos de *Vitex cymosa* Bertero. Rev. Bras. Plantas Medicinais, Botucatu, 8(4):87-91.
- García-García R, López-Malo A, Palou E (2011). Bactericidal action of binary and ternary mixtures of carvacrol, thymol, and eugenol against *Listeria innocua*. J. Food Sci. 76(2):95-100.
- Geromini KVN, Roratto FB, Ferreira FG, Camilotti J, Vidigal TMA, Valle JS, Colauto NB, Linde GA (2015). Fungicidal effect of *Lippia alba* essential oil on a white-rot fungus. Maderas, Cienc. tecnol. 2015 17(1):29-38.
- Glamočlija J, Soković M, Tešević V, Linde G, Colauto NB (2011). Chemical characterization of *Lippia alba* essential oil: an alternative to control green molds. Braz. J. Microbiol. 42:1537-1546.
- Hada D, Sharma K (2014). assessment of antifungal activity of cassia fistula l. fruit pulp against *alternaria solani*. Int. J. Pharm. Pharm. Sci. 7(1):438-441.
- Huang Y, Zhao J, Zhou L, Wang J, Gong Y, Chen X, Guo Z, Wang Q, Jiang W (2010). Antifungal Activity of the Essential Oil of *Illicium verum* Fruit and Its Main Component *trans*-Anethole. Mol. 15:7558-7569.
- Hyldgaard M, Mygind T, Meyer RL (2012). Essential Oils in Food Preservation: Mode of Action, Synergies, and Interactions with Food Matrix Components. Front. Microbiol. 3(12):1-24.
- Ibrahim FAA, Al-Ebady N (2014). Evaluation of Antifungal Activity of Some Plant Extracts and their Applicability in Extending the Shelf Life of Stored Tomato Fruits. J. Food Process. Technol. 5:6.
- Kishore GK, Pande S (2007). Evaluation of essential oils and their components for broad-spectrum antifungal activity and control of late leaf spot and crown rot diseases in peanut. Plant Dis. 91:375-379.
- Knaak N, Fiuza LM (2010). Potencial of essential plant oils to control insects and microorganisms. Neotrop. Biol. Conserv. 5(2):120-132.
- Kurozawa C, Pavan MA (2005). Doenças do tomateiro (*Lycopersicon esculentum* Mill.). In: Kimati H, Amorim L, Rezende JAM, Bergamim FA, Camargo LEA. Manual de Fitopatologia: Doenças das plantas cultivadas, 4. ed. São Paulo SP. Ed. Agron. Ceres. 2:607-626.
- Lee YS, Kim J, Shin SC, Lee SG, Park IK (2008). Antifungal activity of Myrtaceae essential oils and their components against three phytopathogenic fungi. Flavour Fragr. J. 23:23-28.
- Leite GLD, Costa CA, Maia-Almeida CI, Picanço M (2003). Efeito da adubação sobre incidência de traça-do-tomateiro e *alternaria* em plantas de tomate. Hortic. Bras. 21(3):448-451.
- Maia JGS, Silva MD, Luz AI, Zoghbi MDG, Ramos LS (1987). Espécies de Piper da Amazônia ricas em safról. Química Nova 10(3):200-204.
- Ming LC, Maia-Almeida CI, Marques MOM, Conceição DM, Yuhara TY, Leonel S, Tavares RDC, Silva J (2012). The effect of bordaleia and sulfocálcica syrups in different concentrations and ages of plants on leaf diseases of sweet passion fruit in organic cropping. J. Biotechnol. Biodivers. 3(2):30-35.
- Mishra AK, Dubey NK (1994). Evaluation of some essential oils for their toxicity against fungi causing deterioration of stored food commodities. Ap. Environ. Microb. 60:1101-1105.
- Nashwa SM, Abo-Elyousr KA (2012). Evaluation of various plant extracts against the early blight disease of tomato plants under greenhouse and field conditions. Plant Prot. Sci. 48(2):74-79.
- Neely D (1978). Laboratory and greenhouse procedures methods for evaluation fungicides, nematicides and bactericides. Minnesota: An. Phytopathol. Soc. P 140.
- Negrelle RRB, Gomes EC (2007). *Cymbopogon citratus* (DC.) Stapf: chemical composition and biological activities. Rev. Bras. Plantas Med. 9(1):80-92.
- Onawunmi GO, Yisak WA, Ogunlana GO (1984). Antibacterial constituents in the essential oil of *Cymbopogon citratus* (D.C.) Stapf. J. Ethnopharmacol. 12(3):279-286.
- Paes JL, Faroni LRDA, Dhingra OD, Cecon PR, Silva TA (2012). Insecticidal fumigant action of mustard essential oil against *Sitophilus zeamais* in maize grains. Crop. Prot. 34:56-58.
- Pawar VC, Thaker VS (2007). Evaluation of the anti-*Fusarium oxysporum* f. sp. *Cicer* and Anti-*Alternaria porri* effects of some essential oils. World. J. Microb. Technol. 23:1099-1106.
- Perini VDM, Castro HG, dos Santos GR, Chagas Júnior AF, Cardoso

- DP, Aguiar RDS, Soares ADA (2013). Effect of vegetal extract in the inhibition of mycelial growth of *Pyricularia grisea*. J. Biotechnol. Biodivers. 4(1):70-77.
- Salamci E, Kordali S, Kotan R, Cakir A, Kaya Y (2007). Chemical compositions, antimicrobial and herbicidal effects of essential oils isolated from Turkish *Tanacetum aucheranum* and *Tanacetum chiliophyllum* var. *chiliophyllum*. Biochem. Syst. Ecol. 35(9):569-581.
- Salgado APSP, Cardoso MG, Souza PE, Souza JA, Abreu CMP, Simões CMO, Spitzer V (2000). Óleos voláteis. In: Simões C M O et al. Farmacognosia: da planta ao medicamento. 2. ed. Porto Alegre: Universidade Federal do Rio Grande do Sul, Universidade Federal de Santa Catarina. pp. 394-412.
- Sallam NMA (2011). Controlo f tomato Early Blight disease by certain aqueous plant extract. Plant Pathol. J. 10(4):187-191.
- Santana HT, Trindade FTT, Stabeli RG, Silva AAE, Militão JSLT, Facundo VA (2015). Essential oils of leaves of *Piper* species display larvicidal activity against the dengue vector, *Aedes aegypti* (Diptera: Culicidae). Rev. Bras. Plant Med. Campinas 17(1):105-111.
- Souza EL, Lima EO, Freire KRL, Sousa CP (2005). Inhibitory Action of Some Essential Oils and Phytochemicals on the Growth of Various Moulds Isolated From Foods. Braz. Arch. Biol. Technol. 48(2):245-250.
- Souza SMC, Pereira MC, Angélico CLE, Pimenta CJ (2004). Avaliação de óleos essenciais de condimentos sobre o desenvolvimento micelial de fungos associados a produtos de panificação. Ciênc. Agrotéc. 28(3):685-690.
- Soylu EM, Kurt S, Soyulu S (2010). In vitro and in vivo antifungal activities of the essential oils of various plants against tomato grey mould disease agent *Botrytis cinerea*. Int. J. Food Microbiol. 143:183-189.
- Yanar Y, Gökçe A, Kadioglu I, Çam H, Whalon M (2011). *In vitro* antifungal evaluation of various plant extracts against early blight disease (*Alternaria solani*) of potato. Afr. J. Biotechnol. 10(42):8291-8295.
- Zanella CS, Gavassoni WL, Bacchi LMA, Formagio ASN (2015). Activity of plant extracts on the carpogenic germination and mycelial growth of *Sclerotinia sclerotiorum*. Arq. Inst. Biol. São Paulo 82:1-8.